arXiv:1212.6589v2 [quant-ph] 11 Jan 2013

Fluctuation theorems for quantum processes

Tameem Albash,[1, 2] Daniel A. Lidar,[1, 2, 3, 4] Milad Marvian,[2, 3] and Paolo Zanardi[1, 2]
[1]*Department of Physics and Astronomy*
[2]*Center for Quantum Information Science & Technology*
[3]*Department of Electrical Engineering*
[4]*Department of Chemistry*
University of Southern California, Los Angeles, California 90089, USA

We present fluctuation theorems and moment generating function equalities for generalized thermodynamic observables and quantum dynamics described by completely positive trace preserving (CPTP) maps, with and without feedback control. Our results include the quantum Jarzynski equality and Crooks fluctuation theorem, and clarify the special role played by the thermodynamic work and thermal equilibrium states in previous studies. We show that for projective measurements unitality replaces micro-reversibility as the condition for the physicality of the reverse process in our fluctuation theorems. We present an experimental application of our theory to the problem of extracting the system-bath coupling magnitude, which we do for a system of pairs of coupled superconducting flux qubits undergoing quantum annealing.

Introduction.—Fluctuation theorems provide powerful analytical tools for nonequilibrium physics. Some 15 years ago Jarzynski discovered an equality for classical processes that shows how to determine free energy changes by measuring only the work performed on the system, without the need to determine the accompanying entropy changes, and in particular without the requirement that the processes be quasistatic. The classical Jarzynski equality (CJE) [1]

$$\langle e^{-\beta(w-\Delta F)}\rangle = \gamma \ , \tag{1}$$

where $\gamma = 1$, relates the statistical average $\langle\,\rangle$ of the work w done on a driven system to the free energy difference ΔF of the final equilibrium state (whether this state is reached or not) and the initial thermal equilibrium (Gibbs) state, at fixed inverse temperature β. In the presence of feedback control ("Maxwell's demon") the efficacy parameter γ differs from unity, and characterizes the efficacy of feedback and the amount of information extracted [2]. The CJE follows directly from the Tasaki-Crooks fluctuation theorem [3, 4], which relates the probability density function (PDF) of work done in the forward process $P_F(w)$ to the PDF of a reverse process $P_R(w)$:

$$P_F(w)e^{-\beta(w-\Delta F)} = P_R(-w) \ . \tag{2}$$

Many generalizations have been developed (for reviews see, e.g., [5–7]). In particular, the classical results (see also [8]) have been quantized, first for thermal states undergoing unitary evolution [4], and subsequently for thermal states undergoing non-unitary, open system dynamics [9–17], including continuous monitoring [18] and quantum feedback [2, 19, 20].

Here we aim to show that there exists a single unified framework from which all these results can be derived as well as generalized, using only basic tools of the theory of open quantum systems [21] and quantum information theory [22]. To this end we derive a general fluctuation theorem for quantum processes described by completely positive trace-preserving (CPTP) maps $\mathcal{E}$, with or without feedback. Our strategy for doing so leads to a general and simple recipe for writing down fluctuation theorems, not all of which must correspond to a measurable thermodynamic observable or involve a physical reverse process. We show that in order for a PDF for the reverse process to exist, the map $\mathcal{E}$ must be *unital*. In this sense unital channels emerge as playing a crucial role in fluctuation theory, replacing the standard thermodynamic notion of micro-reversibility. Our work illuminates the special role played by the Gibbs state and work measurements, both of which feature prominently in the literature on fluctuation theorems.

We test and confirm our theory via an experiment involving pairs of superconducting flux qubits on a programmable chip, allowing us to empirically determine the first moment of our integral fluctuation theorem. We show that these experimental results can be well explained using a time-dependent Markovian master equation with a free adjustable parameter determining the system-bath coupling strength. As a novel application, our theory allows us to determine this parameter by fitting to the experimental data. We thus establish quantum fluctuation theorems as important tools for studying open quantum systems.

General Fluctuation Theorem.—Consider a fiducial initial state ρ, two sets of generalized measurements $\mathcal{P} := \{P_\alpha\}$ and $\mathcal{Q} := \{Q_\beta\}$ (see also [20]), a CPTP map $\mathcal{E}$ (see also [23]), and a fixed, yet arbitrary distribution $q := \{q_\beta\}$. The quintuple $(\rho, \mathcal{P}, \mathcal{E}, \mathcal{Q}, q)$ is the basic input data describing the problem. The measurement operators P_α satisfy $\sum_\alpha P_\alpha^\dagger P_\alpha = \mathbb{1}$ (the identity operator), and similarly for $\mathcal{Q}$. The CPTP map $\mathcal{E}$ has Kraus operators $\{A_i\}$, i.e., $\mathcal{E}(X) = \sum_i A_i X A_i^\dagger$, where $\sum_i A_i^\dagger A_i = \mathbb{1}$ [24].

We first consider the *forward* process depicted in



FIG. 1. The forward process protocol. A quantum state ρ is prepared, measured ($\mathcal{P}$), evolved via a CPTP map $\mathcal{E}$, and measured again ($\mathcal{Q}$). The final state ρ_f thus obtained is in general different from the virtual final state ρ_q.

Fig. 1. A mixed state ensemble $\rho_p = \sum_\alpha p_\alpha \rho_\alpha$ is prepared by measuring $\mathcal{P}$, so that the normalized state $\rho_\alpha = P_\alpha \rho P_\alpha^\dagger / p_\alpha$ has probability $p_\alpha = \mathrm{Tr}[P_\alpha^\dagger P_\alpha \rho]$. Next ρ_α evolves under $\mathcal{E}$, and finally the measurement $\mathcal{Q}$ is performed. The conditional probability of observing outcome β given outcome α is then $p_{\beta|\alpha} = \mathrm{Tr}[Q_\beta^\dagger Q_\beta \mathcal{E}(\rho_\alpha)]$. The marginal probability distribution of outcomes is $f := \{f_\beta\}$, where $f_\beta = \sum_\alpha p_{\beta|\alpha} p_\alpha = \sum_\alpha p_{(\alpha,\beta)}$, and where $p_{(\alpha,\beta)}$ is the joint probability distribution.

We next define the (virtual) final state $\rho_q := \sum_{\beta=1}^d q_\beta \sigma_\beta$, where $\sigma_\beta := Q_\beta^\dagger Q_\beta > 0$ is a normalized state having probability $q_\beta > 0 \ \forall \beta$. Note that $d = \mathrm{Tr}[\mathbb{1}]$ is the system Hilbert space dimension and that $f_\beta = \mathrm{Tr}\left[\sigma_\beta \mathcal{E}(\rho_p)\right]$. Let the random variable V be a real-valued function parametrized by the measurement outcomes $\{\alpha, \beta\}$. V will play the role of a generalized thermodynamic observable [25]. Its PDF is

$$P_\mathcal{E}(v) := \sum_{\alpha,\beta} \delta(v - V_{\alpha\beta}) p_{(\alpha,\beta)} \tag{3a}$$

$$= \sum_{\alpha,\beta} \delta(v - V_{\alpha\beta}) \mathrm{Tr}[\sigma_\beta \mathcal{E}(\rho_\alpha)] p_\alpha \ . \tag{3b}$$

Let us now choose $V_{\alpha\beta} = \ln(p_\alpha / q_\beta)$ (this choice is by no means unique [26]), and also define the generalized reverse thermodynamic observable $\tilde{V}_{\beta\alpha} := -V_{\alpha\beta} = \ln(q_\beta / p_\alpha)$. Then, using the Dirac delta properties $\delta(x - x_0) f(x) = \delta(x - x_0) f(x_0)$, and $\delta(x) = \delta(-x)$, we find

$$P_\mathcal{E}(v) e^{-v} = \sum_{\alpha,\beta} \delta(v - V_{\alpha\beta}) \mathrm{Tr}[\sigma_\beta \mathcal{E}(\rho_\alpha)] q_\beta \tag{4a}$$

$$= \sum_{\alpha,\beta} \delta(-v - \tilde{V}_{\beta\alpha}) \mathrm{Tr}[\rho_\alpha \mathcal{E}^*(\sigma_\beta)] q_\beta \ , \tag{4b}$$

where we used the dual map $\mathcal{E}^*$ with Kraus operators $\{A_i^\dagger\}$, i.e., $\mathcal{E}^*(X) = \sum_i A_i^\dagger X A_i$, for which $\mathrm{Tr}[A\mathcal{E}(B)] = \mathrm{Tr}[B\mathcal{E}^*(A)]$ for any pair of operators A and B.

Comparing Eqs. (3b) and (4b) we see that the latter can be identified with $\tilde{P}_{\mathcal{E}^*}(-v)$. We have therefore arrived at our fluctuation theorem for CPTP maps:

$$P_\mathcal{E}(v) e^{-v} = \tilde{P}_{\mathcal{E}^*}(-v) \ . \tag{5}$$

Note the obvious similarity to the Tasaki-Crooks fluctuation theorem, Eq. (2).

We define the *efficacy* γ [2, 23] as

$$\gamma := \int_{-\infty}^{\infty} dv \, \tilde{P}_{\mathcal{E}^*}(v) = \sum_\alpha \mathrm{Tr}[\rho_q \mathcal{E}(\rho_\alpha)] = \sum_\alpha \mathrm{Tr}[\rho_\alpha \mathcal{E}^*(\rho_q)] \ . \tag{6}$$

Note that $\tilde{P}_{\mathcal{E}^*}$ need not be a probability distribution (it is not necessarily normalized). Upon integration, our detailed fluctuation theorem gives an integral fluctuation theorem, which we call the quantum Jarzynski equality (QJE), as it generalizes the CJE, Eq. (1):

$$\langle e^{-v} \rangle = \int_{-\infty}^{\infty} dv \, P_\mathcal{E}(v) e^{-v} = \gamma \ . \tag{7}$$

Using Jensen's inequality we have $\langle e^f \rangle \geq e^{\langle f \rangle}$ and thus find a generalized 2nd law of thermodynamics (we clarify this claim below) [27]:

$$\langle v \rangle \geq -\ln \gamma \ . \tag{8}$$

We can substantially generalize the QJE Eq. (7) in terms of the moment generating functions for the map $\mathcal{E}$ and its dual,

$$\chi_\mathcal{E}(\lambda) := \int_{-\infty}^{\infty} dv P_\mathcal{E}(v) e^{\lambda v} \ , \ \tilde{\chi}_{\mathcal{E}^*}(\lambda) := \int_{-\infty}^{\infty} dv \tilde{P}_{\mathcal{E}^*}(v) e^{\lambda v} . \tag{9}$$

We then find after some calculation [28] our central result:

$$\chi_\mathcal{E}(\lambda - 1) = \tilde{\chi}_{\mathcal{E}^*}(-\lambda) \ , \tag{10}$$

It extends the integral fluctuation relation Eq. (7) to all moments of the PDF $P_\mathcal{E}(v)$. For example, setting $\lambda = 0$, we recover the QJE Eq. (7): $\chi_\mathcal{E}(-1) = \langle e^{-v} \rangle = \tilde{\chi}_{\mathcal{E}^*}(0) = \gamma$.

Moreover, using $\langle v \rangle = \frac{d}{d\lambda} \chi_\mathcal{E}(\lambda)\big|_{\lambda=0}$ we find [28]

$$\langle v \rangle = H(f \| q) + H(f) - H(p) \ , \tag{11}$$

where $H(f \| q) = -\sum_\beta f_\beta \ln(q_\beta) - H(f)$ is the relative entropy (Kullback-Leibler divergence), and $H(f) = -\sum_\beta f_\beta \ln(f_\beta)$ is the Shannon entropy.

Projective measurements.—An important special case arises when the measurements $\mathcal{P}$ and $\mathcal{Q}$ are projective and when moreover we prepare $\rho_p = \sum_\alpha p_\alpha P_\alpha$ and let $\rho_q = \sum_\beta q_\beta Q_\beta$, with $\{P_\alpha\}$ and $\{Q_\beta\}$ rank-1 projectors (pure states). For $V_{\alpha\beta} = \ln(p_\alpha / q_\beta)$, we obtain $\gamma = \mathrm{Tr}\left[\rho_q \mathcal{E}(\mathbb{1})\right] = \mathrm{Tr}\left[\mathcal{E}^*(\rho_q)\right]$ (see also [23]). In this case, $\tilde{P}_{\mathcal{E}^*}$ is a probability distribution if $\mathcal{E}^*$ is itself a CP map, i.e., $\mathcal{E}$ is unital [$\mathcal{E}(\mathbb{1}) = \mathbb{1}$] . If it is unital then $\gamma = 1$ [29], and we can interpret $\tilde{P}_{\mathcal{E}^*}(v)$ as the probability density associated with the following *reverse* process: i) prepare the state $\rho_q = \sum_\beta q_\beta Q_\beta$, ii) measure $\mathcal{Q}$, iii) evolve via $\mathcal{E}^*$, iv) measure $\mathcal{P}$. *Thus unitality replaces micro-reversibility.* Why are unital channels singled out? Whereas classical micro-reversibility imposes a specific relation between forward and time-reversed phase space


information-geometric distance
(2)
$\mathcal{E}\left(\rho_p\right)$
ρ_q
Von Neumann entropy change
physical process
(1)
(3)
ρ_p
Von Neumann entropy change
virtual process

FIG. 2. Distance-entropy diagram illustrating the generalized 2nd law of thermodynamics (see text for details).

paths [3], a unique and suggestive feature of unital channels is that they can always be written as an affine combination of reversible channels, i.e., $\mathcal{E}(\rho)=\sum_j u_j U_j \rho U_j^\dagger$, where U_j is unitary, $u_j \in \mathbb{R}$ and $\sum_j u_j = 1$ [30].

Using Eq. (9), we then obtain [28]

$$\chi_{\mathcal{E}}(\lambda)=\operatorname{Tr}\left[\rho_q^{-\lambda}\mathcal{E}\left(\rho_p^{\lambda+1}\right)\right] \ , \tag{12}$$

and consequently:

$$\langle v\rangle = S(\mathcal{E}(\rho_p)) - S(\rho_p) + S(\mathcal{E}(\rho_p)\|\rho_q) \tag{13a}$$

$$= S(\rho_q) - S(\rho_p) + \operatorname{Tr}[(\rho_q - \mathcal{E}(\rho_p))\ln(\rho_q)] \ , \tag{13b}$$

where the quantum relative entropy $S\left(\rho\|\sigma\right) = -\operatorname{Tr}[\rho\ln\sigma] - S(\rho)$, and $S(\rho) = -\operatorname{Tr}[\rho\ln\rho]$ is the von Neumann entropy. This generalizes the result for the mean irreversible entropy production of Refs. [31, 32]. If $S(\mathcal{E}(\rho_p)) = S(\rho_p)$ (e.g., when $\mathcal{E}$ is unitary) then $\langle v\rangle = S\left(\mathcal{E}(\rho_p)\|\rho_q\right)$. If ρ_q is $\mathcal{E}$-invariant $[\mathcal{E}(\rho_q) = \rho_q]$ and $\mathcal{E}$ is unitary (as in a quantum quench), then $\langle v\rangle = S\left(\rho_p\|\rho_q\right)$. If $\mathcal{E}(\rho_p) = \rho_q$ then $\langle v\rangle = S(\rho_q) - S(\rho_p)$ [33].

Eq. (13) has an interesting interpretation, illustrated in Fig. 2. Referring first to Eq. (13a), side (1) of the triangle represents the von Neumann entropy change $(1) := S(\mathcal{E}(\rho_p)) - S(\rho_p)$ occurring in the physical process enacted by $\mathcal{E}$, while side (2) is an information-geometric measure of the distance $(2) := S(\mathcal{E}(\rho_p)\|\rho_q)$ between the evolved state $\mathcal{E}(\rho_p)$ and the "virtual" one ρ_q [34]. On the other hand, referring to Eq. (13b), side (2) can also represent $(2') := \operatorname{Tr}[(\rho_q - \mathcal{E}(\rho_p))\ln(\rho_q)] \leq \|\mathcal{E}(\rho_p) - \rho_q\|_1 |\ln(\min_\beta q_\beta)|$, which is again related to the information-theoretic distance between the evolved and virtual state. Side (3) represent the von Neumann entropy change, i.e., $(3) := S(\rho_q) - S(\rho_p)$.

In the thermal case $\rho_p = e^{-\beta H_i}/Z_i$ and $\rho_q = e^{-\beta H_f}/Z_f$ (where $H_{i/f}$ is the initial/final Hamiltonian), we find $(2') = \beta(\operatorname{Tr}[H_f\mathcal{E}(\rho_p)] - \operatorname{Tr}[H_f\rho_q]) =: -\beta Q$. Here Q represents the *heat exchange* in the (virtual) undriven relaxation between the evolved and virtual states, $\mathcal{E}(\rho_p) \mapsto \rho_q$. In the thermal case (3) amounts to a *thermodynamical entropy* change ΔS_{qp}. Thus in this case, using Eq. (8) and assuming unitality ($\gamma = 1$), we have the 2nd law in the Clausius form $\Delta S_{qp} \geq \beta Q$. This clarifies why Eq. (8) can be interpreted as a generalized 2nd law of thermodynamics.

Including feedback.— Suppose we repeat the forward protocol of Fig. 1 in the projective measurement case, but denote the CPTP map by $\bar{\mathcal{E}}$ and the final measurement by $\mathcal{Q} = \{Q_j\}$. Depending on the measurement outcome j, we apply an additional CP map $\bar{\mathcal{E}}_j$ to the resulting state Q_j. This constitutes a *feedback* step, and generalizes earlier work which considered only unitary feedback maps [2, 19, 20]. Next we apply another projective measurement $\mathcal{Q}'_j = \{Q_{\beta|j}\}$, labeled by the outcomes β. Thus $\mathcal{E}$ from Fig. 1 is replaced by $\mathcal{E}_j := \bar{\mathcal{E}}_j \circ \mathcal{Q} \circ \bar{\mathcal{E}}$, and $\mathcal{Q}$ by $\mathcal{Q}'_j$. Given an initial state P_α, the probability of observing outcomes j and β is $p_{(j,\beta)|\alpha} = \operatorname{Tr}\left[Q_{\beta|j}\mathcal{E}_j(P_\alpha)\right]$, and the joint distribution is $p_{(\alpha,j,\beta)} = p_{(j,\beta)|\alpha}p_\alpha$. As in the feedback-free case, we can construct the PDF associated with the generalized thermodynamic observable V as:

$$P_{\mathcal{E}} := \sum_{\alpha,j,\beta}\delta(v - V_{\alpha j\beta})p_{(\alpha,j,\beta)} \tag{14a}$$

$$= \sum_{\alpha,j,\beta}\delta(v - V_{\alpha j\beta})\operatorname{Tr}[Q_{\beta|j}\mathcal{E}_j(P_\alpha)]p_\alpha \ , \tag{14b}$$

where the notation $P_{\mathcal{E}}$ $(\tilde{P}_{\mathcal{E}^*})$ is shorthand for $P_{\{\mathcal{E}_j\}}$ $(\tilde{P}_{\{\mathcal{E}_j^*\}})$. We now choose $V_{\alpha j\beta} = \ln\left(p_\alpha/q_{\beta|j}\right) = -\tilde{V}_{\beta j\alpha}$, where $q_j := \{q_{\beta|j}\}$ is an arbitrary, fixed distribution, associated with the (virtual) state $\rho_{q|j} := \sum_\beta q_{\beta|j}Q_{\beta|j}$. Then

$$P_{\mathcal{E}}(v)e^{-v} = \sum_{\alpha,j,\beta}\delta(-v - \tilde{V}_{\beta j\alpha})\operatorname{Tr}[P_\alpha\mathcal{E}_j^*(Q_{\beta|j})]q_{\beta|j} \tag{15a}$$

$$=: \tilde{P}_{\mathcal{E}^*}(-v) \ . \tag{15b}$$

Integrating, we find a generalized integral fluctuation theorem in the presence of feedback:

$$\langle e^{-v}\rangle = \sum_j \operatorname{Tr}[\rho_{q|j}\bar{\mathcal{E}}_j(\mathbb{1})] = \sum_j \operatorname{Tr}[\mathcal{E}_j^*\left(\rho_{q|j}\right)] =: \gamma \ . \tag{16}$$

Generalizing Eqs. (10) and (12), we find the moment generating function for the feedback case

$$\chi_{\mathcal{E}}(\lambda - 1) = \sum_j \operatorname{Tr}\left[\rho_{q|j}^{1-\lambda}\mathcal{E}_j\left(\rho_p^\lambda\right)\right] = \tilde{\chi}_{\mathcal{E}^*}(-\lambda) \ . \tag{17}$$

We show in [28] how these results allow us to recover known quantum fluctuation theorems with feedback [35].

Experimental application of the open-system QJE.— Compared to the classical case (e.g., [36–41]), there has been relatively little work on experimental tests or applications of the quantum version of the Jarzysnki-Crooks relations [42]. We now present an application of our generalized QJE [Eq. (7)] to the problem of extracting the system-bath coupling magnitude, using both numerical simulations and an experiment with a quantum annealing chip comprising superconducting flux qubits (for experimental details see [28]). The qubits are described by the tranverse-field Ising Hamiltonian $H_S(t) = -A(t)\sum_i \sigma_i^x + B(t)H_{\text{Ising}}$, where $H_{\text{Ising}} = -\sum_{i=1}^N h_i\sigma_i^z - \sum_{i<j}^N J_{ij}\sigma_i^z\sigma_j^z$, and where the σ's are standard Pauli operators. The magnetic fields h_i and couplings (superconducting inductances) J_{ij} are programmable. This open system can be described by an adiabatic Markovian master equation [43] derived in Ref. [44] (see [28] for essential details). The annealing functions $A(t)$ and $B(t)$ satisfy $A(0), B(t_f) \neq 0$, $A(t_f), B(0) = 0$, where t_f is the total annealing time. The CPTP map $\mathcal{E}$ is the one generated by the master equation and is *not* unital [28]. We consider projective measurements that prepare $p_\alpha = e^{-\beta\varepsilon_\alpha(0)}/Z(0)$, $q_\beta = e^{-\beta\varepsilon_\beta(t_f)}/Z(t_f)$, where $\varepsilon_{\alpha,\beta}(t)$ are the instantaneous eigenenergies of $H_S(t)$, so that our generalized thermodynamic observable $V_{\alpha\beta} = \ln(p_\alpha/q_\beta)$ is given by $V_{\alpha\beta} = \beta(\varepsilon_\beta(t_f) - \varepsilon_\alpha(0)) - \beta(F(t_f) - F(0))$, where the free energy $F = -\ln(Z)/\beta$. Note that for an open quantum system, $\varepsilon_\beta(t_f) - \varepsilon_\alpha(0)$ does *not* correspond to the work done on the system. Equations (6) and (7) yield

$$\langle e^{-\beta(\Delta E - \Delta F)}\rangle = \gamma = \mathrm{Tr}\left[\mathcal{E}^*(\rho_{\mathrm{G}}(t_f))\right] \ , \tag{18}$$

where $\rho_{\mathrm{G}}(t) = \exp(-\beta H_S(t))/Z(t)$ denotes the Gibbs state associated with $H_S(t)$, with $\rho_p \equiv \rho_{\mathrm{G}}(0)$ and $\rho_q \equiv \rho_{\mathrm{G}}(t_f)$, ΔE is a random variable taking values in the set $\{\varepsilon_\beta(t_f) - \varepsilon_\alpha(0)\}$, and $\Delta F := F(t_f) - F(0)$. Equation (13a) gives:

$$\begin{aligned}\beta\left(\langle\Delta E\rangle - \Delta F\right) = S\left(\rho(t_f)\|\rho_{\mathrm{G}}(t_f)\right) + S(\rho(t_f))\\ - S\left(\rho_{\mathrm{G}}(0)\right) \ ,\end{aligned} \tag{19}$$

where we have denoted $\rho(t_f) = \mathcal{E}(\rho_{\mathrm{G}}(0))$.

For concreteness we consider a two-qubit system with $H_{\text{Ising}} = -\frac{1}{3}\sum_{i=1}^2 \sigma_i^z - J\sigma_1^z\sigma_2^z$. We checked the equality expressed in Eq. (18) by independently numerically simulating its two sides for this model using our master equation, and find essentially perfect agreement. The same holds for Eq. (13a) [28]. We tested the same two-qubit system experimentally. The system was initialized in the Gibbs state (at $T = 17$mK) and we measured the occupancy of each final energy eigenstate (of H_{Ising}), repeating the quantum annealing process thousands of times. The empirical relative occupancy corresponds directly to the probability f_β of measuring energy $\varepsilon_\beta(t_f)$. Therefore, this allows us to experimentally determine $\langle\varepsilon(t_f)\rangle = \sum_\beta \varepsilon_\beta(t_f) f_\beta$. We also know $\langle\varepsilon(0)\rangle$ from the initial Gibbs state and the value of $A(0)$. We thus determine

$$\begin{aligned}\langle v\rangle &= \beta\sum_{\alpha,\beta}\left(\varepsilon_\beta(t_f) - \varepsilon_\alpha(0) - \Delta F\right)p_{(\alpha,\beta)}\\ &= \beta\left(\langle\varepsilon(t_f)\rangle - \langle\varepsilon(0)\rangle - \Delta F\right) \ ,\end{aligned} \tag{20}$$



FIG. 3. Experimental results (blue dots with error bars) for $\langle v\rangle$ [from Eq. (20)] and the best fit using the adiabatic Markovian master equation (red ×) with the extracted value $\kappa = 1.83\times10^{-3}$. Main panel: as a function of J, with $t_f = 5\mu$s in both the numerical simulations and the experiment. Inset: as a function of t_f, with $J = 1/2$ in both the numerical simulations and the experiment.

where we compute ΔF using exact diagonalization [28]. We show these results in Fig. 3 as a function of t_f and ferromagnetic coupling strength J. Our master equation has two free parameters: the high-frequency cut-off ω_c, which we set to 8π [44], and the system-bath coupling magnitude $\kappa = g^2\eta/\hbar^2$, where g is the system-bath coupling constant and η characterizes the Ohmic bath [28]. Remarkably, by simultaneously minimizing the deviation between the numerical solution of our master equation and the experimental data for $\langle v\rangle$ as a function of J and t_f allowed us to extract the system-bath coupling magnitude κ from Eq. (20) [28].

Why does $\langle v\rangle$ display a minimum as a function of J (main panel of Fig. 3)? In our experiment $\beta A(0) \sim \beta B(t_f) \approx 15$, so that the Gibbs state is almost pure, i.e., both $S(\rho_{\mathrm{G}}(0))$ and $S(\rho_{\mathrm{G}}(t_f)) \ll 1$. Therefore *we are effectively measuring the information-theoretic distance* $S\left(\rho(t_f)\|\rho_{\mathrm{G}}(t_f)\right)$. Increasing J at fixed temperature $kT = 1/\beta$ is like decreasing T while fixing J. Thus the system requires more time to equilibrate as J grows, but we keep t_f fixed. On the other hand, as J becomes very small the ground and first excited states become degenerate, so the excitation probability increases, and the system is again farther from equilibrium. Also, as we increase the annealing time, $\rho(t_f)$ becomes closer to the Gibbs state, causing $S\left(\rho(t_f)\|\rho_{\mathrm{G}}(t_f)\right)$ to decrease as observed in the inset.

Conclusions.— We presented fluctuation theorems and moment generating functions for CPTP maps, thus generalizing previous work on the Jarzynski-Crooks relations and the 2nd law for open quantum systems, including processes with feedback. We performed an experiment using superconducting flux qubits that matches the fluctuation theorem protocols, and used this experiment to extract the system-bath coupling for an adiabatic Markovian master equation that nicely matches the experimental results. Our work ties together key ideas from statistical mechanics, quantum information theory, and the theory of open quantum systems, and paves the way to experimental tests and applications of fluctuation theorems in the most general setting of open quantum system dynamics.

We thank S. Deffner and P. Talkner for useful comments. This research was supported by the ARO MURI grant W911NF-11-1-0268, the Lockheed Martin Corporation, and by NSF grant numbers PHY- 969969 and PHY-803304 (to P.Z. and D.A.L.).

[1] C. Jarzynski, Phys. Rev. Lett. **78**, 2690 (1997).

[2] T. Sagawa and M. Ueda, Phys. Rev. Lett. **104**, 090602 (2010).

[3] G. E. Crooks, Phys. Rev. E **60**, 2721 (1999).

[4] H. Tasaki, (2000), arXiv:cond-mat/0009244.

[5] C. Bustamante, J. Liphardt, and F. Ritort, Phys. Today **58**, 43 (2005).

[6] C. Jarzynski, Annu. Rev. Cond. Mat. Phys. **2**, 329 (2011).

[7] M. Campisi, P. Hänggi, and P. Talkner, Rev. Mod. Phys. **83**, 771 (2011).

[8] U. Seifert, Phys. Rev. Lett. **95**, 040602 (2005).

[9] S. Yukawa, J. Phys. Soc. Jpn **69**, 2367 (2000).

[10] S. Mukamel, Phys. Rev. Lett. **90**, 170604 (2003).

[11] T. Monnai, Phys. Rev. E **72**, 027102 (2005).

[12] P. Talkner, E. Lutz, and P. Hänggi, Phys. Rev. E **75**, 050102 (2007).

[13] G. E. Crooks, J. Stat. Mech. Theor. Exp. **2008**, P10023 (2008).

[14] H. T. Quan and H. Dong, arXiv:0812.4955.

[15] M. Campisi, P. Talkner, and P. Hänggi, Phys. Rev. Lett. **102**, 210401 (2009).

[16] P. Talkner, M. Campisi, and P. Hänggi, J. Stat. Mech. Theor. Exp. **2**, 25 (2009).

[17] S. Deffner and E. Lutz, Phys. Rev. Lett. **107**, 140404 (2011).

[18] M. Campisi, P. Talkner, and P. Hänggi, Phys. Rev. Lett. **105**, 140601 (2010).

[19] T. Sagawa and M. Ueda, Phys. Rev. Lett. **100**, 080403 (2008).

[20] Y. Morikuni and H. Tasaki, J. Stat. Phys. **143**, 1 (2011).

[21] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, Oxford, 2002).

[22] M. Nielsen and I. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, England, 2000).

[23] D. Kafri and S. Deffner, Phys. Rev. A **86**, 044302 (2012).

[24] Note that generalized measurements include projective measurements, positive operator valued measures (POVM), and weak measurements as special cases [22, 45].

[25] Note that work is not a quantum observable [12], and that we use the term ‘observable’ in a loose sense.

[26] We stress that we are not restricted to the choice $V_{\alpha\beta} = \ln(p_\alpha/q_\beta)$. For example, if we choose $V_{\alpha\beta} = \ln\left(p_{\beta|\alpha}/f_\beta\right)$ then Eq. (3a) yields $P_{\mathcal{E}}(v)e^{-v} = \sum_{\alpha,\beta}\delta(v-V_{\alpha\beta})p_\alpha f_\beta$. In this case Eq. (7) immediately yields the QJE with $\gamma = 1$.

[27] We further remark concerning Eq. (8) that, upon applying $|\mathrm{Tr}[AB]| \leq \|A\|_\infty \|B\|_1$ (the product of the sup-operator and trace norms [46]) to Eq. (6), we find that $\gamma \leq \|\mathcal{E}(\mathbb{1})\|_\infty \leq d$.

[28] See Supplemental Material for additional details and proofs.

[29] The observation that unital channels yield $\gamma = 1$ was first stated in Ref. [47].

[30] For a proof see Theorem 1 in [48]. In the special case when $\{u\}$ is a probability distribution, one can interpret the unital channel as representing the unitary evolution U_j occurring with probability u_j, and its dual as the time-reversed unitary $U_j^\dagger$ occurring with the same probability.

[31] S. Deffner and E. Lutz, Phys. Rev. Lett. **105**, 170402 (2010).

[32] R. Dorner, J. Goold, C. Cormick, M. Paternostro, and V. Vedral, Phys. Rev. Lett. **109**, 160601 (2012).

[33] Known quantum fluctuation theorems follow from our formalism. For example, we show in [28] how Eq. (13a) reduces to the standard statement of the 2nd law for isothermal processes, $\beta\left(\langle w\rangle - \Delta F\right) = \langle \Delta S_{\mathrm{irr}}\rangle \geq 0$, where $\langle \Delta S_{\mathrm{irr}}\rangle$ is the mean irreversible entropy production [17, 31], after we choose p and q as Gibbs distributions and $\mathcal{E}$ as a unitary evolution. We also note that a calculation of $\langle v^2\rangle = \frac{d^2}{d\lambda^2}\chi_{\mathcal{E}}(\lambda)\Big|_{\lambda=0}$ would yield a generalized fluctuation-dissipation theorem.

[34] The relative entropy is not strictly a distance (since it is not symmetric and does not satisfy the triangle inequality), but a divergence.

[35] Although we have identified $P_{\mathcal{E}}(v)e^{-v} \equiv P_{\{\mathcal{E}_j\}}(v)e^{-v}$ with $\tilde{P}_{\mathcal{E}^*}(-v) \equiv \tilde{P}_{\{\mathcal{E}_j^*\}}(-v)$, it is important to note that $\mathcal{E}^*$ does not coincide with the reverse process because (unlike in the feedback-free case) there is now no unique association between the initial α and final β measurement outcome indices: the same (α,β) pair is connected via different j values of the intermediate measurement outcomes.

[36] G. M. Wang, E. M. Sevick, E. Mittag, D. J. Searles, and D. J. Evans, Phys. Rev. Lett. **89**, 050601 (2002).

[37] D. Collin, F. Ritort, C. Jarzynski, S. Smith, I. Tinoco, and C. Bustamante, Nature **437**, 231 (2005).

[38] I. Junier, A. Mossa, M. Manosas, and F. Ritort, Phys. Rev. Lett. **102**, 070602 (2009).

[39] N. Garnier and S. Ciliberto, Phys. Rev. E **71**, 060101 (2005).

[40] F. Douarche, S. Ciliberto, and A. Petrosyan, Journal of Statistical Mechanics: Theory and Experiment **2005**, P09011 (2005).

[41] S. Toyabe, T. Sagawa, M. Ueda, E. Muneyuki, and M. Sano, Nat. Phys. **6**, 988 (2010).

[42] Existing data on x-ray spectra of simple metals [49], experiments on a driven single qubit (defect center in diamond) [50], and bidirectional single-electron counting [51] have been used to verify the previously derived quantum fluctuation relations.

[43] Fluctuation theorems for Markovian master equations have been studied in Refs. [52, 53].

[44] T. Albash, S. Boixo, D. A. Lidar, and P. Zanardi, New J. of Phys. **14**, 123016 (2012).

[45] O. Oreshkov and T. A. Brun, Phys. Rev. Lett. **95**, 110409 (2005).

[46] D. A. Lidar, P. Zanardi, and K. Khodjasteh, Phys. Rev. A **78**, 012308 (2008).
[47] D. Kafri and S. Deffner, arXiv:1207.2978v1 [quant-ph].
[48] C. Mendl and M. Wolf, Comm. Math. Phys. **289**, 1057 (2009).
[49] M. Heyl and S. Kehrein, Phys. Rev. Lett. **108**, 190601 (2012).
[50] S. Schuler, T. Speck, C. Tietz, J. Wrachtrup, and U. Seifert, Phys. Rev. Lett. **94**, 180602 (2005).
[51] Y. Utsumi, D. S. Golubev, M. Marthaler, K. Saito, T. Fujisawa, and G. Schön, Phys. Rev. B **81**, 125331 (2010).
[52] M. Esposito and S. Mukamel, Phys. Rev. E **73**, 046129 (2006).
[53] R. Chetrite and K. Mallick, Journal of Statistical Physics **148**, 480 (2012).
[54] A. J. Leggett, S. Chakravarty, A. T. Dorsey, M. P. A. Fisher, A. Garg, and W. Zwerger, Rev. Mod. Phys. **59**, 1 (1987).
[55] R. Harris, M. W. Johnson, T. Lanting, A. J. Berkley, J. Johansson, P. Bunyk, E. Tolkacheva, E. Ladizinsky, N. Ladizinsky, T. Oh, F. Cioata, I. Perminov, P. Spear, C. Enderud, C. Rich, S. Uchaikin, M. C. Thom, E. M. Chapple, J. Wang, B. Wilson, M. H. S. Amin, N. Dickson, K. Karimi, B. Macready, C. J. S. Truncik, and G. Rose, Phys. Rev. B **82**, 024511 (2010).
[56] A. J. Berkley, M. W. Johnson, P. Bunyk, R. Harris, J. Johansson, T. Lanting, E. Ladizinsky, E. Tolkacheva, M. H. S. Amin, and G. Rose, Superconductor Science and Technology **23**, 105014 (2010).
[57] M. W. Johnson, P. Bunyk, F. Maibaum, E. Tolkacheva, A. J. Berkley, E. M. Chapple, R. Harris, J. Johansson, T. Lanting, I. Perminov, E. Ladizinsky, T. Oh, and G. Rose, Superconductor Science and Technology **23**, 065004 (2010).

Supplemental Material for “Fluctuation theorems for quantum processes”

To distinguish main text equation numbers from equation numbers in this Supplemental Material, we denote equation number x from the main text by “Eq.M. (x)”, while equation numbers cross-referenced in the Supplemental Material are denoted by the usual “Eq. (x)”.

DERIVATION OF EQ.M. (10) AND EQ.M. (12)

Using Eq.M. (3b) and Eq.M. (9) we have

$$\chi_{\mathcal{E}}(\lambda) := \int_{-\infty}^{\infty} dv P_{\mathcal{E}}(v) e^{\lambda v} \tag{1a}$$

$$= \int_{-\infty}^{\infty} dv e^{\lambda v} \sum_{\alpha,\beta} \delta(v - V_{\{\alpha,\beta\}}) \mathrm{Tr}[\sigma_\beta \mathcal{E}(\rho_\alpha)] p_\alpha \tag{1b}$$

$$= \sum_{\alpha,\beta} (p_\alpha / q_\beta)^\lambda \, \mathrm{Tr}\left[\sigma_\beta \mathcal{E}(\rho_\alpha)\right] p_\alpha \tag{1c}$$

$$= \mathrm{Tr}\left[\mathcal{E}\left(\sum_\alpha p_\alpha^{\lambda+1} \rho_\alpha\right) \sum_\beta q_\beta^{-\lambda} \sigma_\beta\right] , \tag{1d}$$

so that

$$\chi_{\mathcal{E}}(\lambda - 1) = \mathrm{Tr}\left[\mathcal{E}\left(\sum_\alpha p_\alpha^{\lambda} \rho_\alpha\right) \sum_\beta q_\beta^{1-\lambda} \sigma_\beta\right] \tag{2a}$$

$$= \mathrm{Tr}\left[\sum_\alpha p_\alpha^{\lambda} \rho_\alpha \, \mathcal{E}^* \left(\sum_\beta q_\beta^{1-\lambda} \sigma_\beta\right)\right] . \tag{2b}$$

On the other hand, using Eq.M. (4b) and Eq.M. (9) we have

$$\tilde{\chi}_{\mathcal{E}^*}(-\lambda) = \int_{-\infty}^{\infty} dv \tilde{P}_{\mathcal{E}^*}(v) e^{-\lambda v} \tag{3a}$$

$$= \int_{-\infty}^{\infty} dv e^{-\lambda v} \sum_{\alpha,\beta} \delta(v - \tilde{V}_{\{\beta,\alpha\}}) \mathrm{Tr}[\rho_\alpha \mathcal{E}^*(\sigma_\beta)] q_\beta \tag{3b}$$

$$= \sum_{\alpha,\beta} (p_\alpha / q_\beta)^\lambda \, \mathrm{Tr}\left[\rho_\alpha \mathcal{E}^*(\sigma_\beta)\right] q_\beta \tag{3c}$$

$$= \mathrm{Tr}\left[\sum_\alpha p_\alpha^{\lambda} \rho_\alpha \, \mathcal{E}^* \left(\sum_\beta q_\beta^{1-\lambda} \sigma_\beta\right)\right] \tag{3d}$$

$$= \chi_{\mathcal{E}}(\lambda - 1) \; . \tag{3e}$$

In the case of projective measurements and and when $\rho_p = \sum_\alpha p_\alpha P_\alpha$, $\rho_q = \sum_\beta q_\beta Q_\beta$, with $\{P_\alpha\}$ and $\{Q_\beta\}$ rank-1 projectors, Eq. (1d) directly becomes Eq.M. (12).

DERIVATION OF THE ENTROPY FORMULAS FOR $\langle v \rangle$

Generalized measurements: Eq.M. (11)

We prove Eq.M. (11) from the first moment expression $\langle v \rangle = \frac{d}{d\lambda} \chi_{\mathcal{E}}(\lambda)\Big|_{\lambda=0}$. Then, using Eq. (1d)

$$\langle v \rangle = \frac{d}{d\lambda} \mathrm{Tr}\left[\mathcal{E}\left(\sum_\alpha p_\alpha^{\lambda+1} \rho_\alpha\right) \sum_\beta q_\beta^{-\lambda} \sigma_\beta\right] \Big|_{\lambda=0} \tag{4a}$$

$$\begin{aligned} = \; & \mathrm{Tr}\left[\mathcal{E}\left(\sum_\alpha p_\alpha^{\lambda+1} \ln(p_\alpha) \rho_\alpha\right) \sum_\beta q_\beta^{-\lambda} \sigma_\beta\right] \Big|_{\lambda=0} \\ & - \mathrm{Tr}\left[\mathcal{E}\left(\sum_\alpha p_\alpha^{\lambda+1} \rho_\alpha\right) \sum_\beta q_\beta^{-\lambda} \ln(q_\beta) \sigma_\beta\right] \Big|_{\lambda=0} \end{aligned} \tag{4b}$$

$$= \mathrm{Tr}\left[\mathcal{E}\left(\sum_\alpha p_\alpha \ln(p_\alpha) \rho_\alpha\right)\right] - \mathrm{Tr}\left[\mathcal{E}(\rho_p) \sum_\beta \ln(q_\beta) \sigma_\beta\right] \tag{4c}$$

$$= \sum_\alpha p_\alpha \ln(p_\alpha) - \sum_\beta f_\beta \ln(q_\beta) \; , \tag{4d}$$

where to arrive at Eq. (4c) we used the fact that $\sum_\beta \sigma_\beta = \sum_\beta Q_\beta^\dagger Q_\beta = \mathbb{1}$, to arrive at Eq. (4d) that $\mathcal{E}$ is trace-preserving and $\mathrm{Tr}[\rho_\alpha] = 1$, and observed that $\mathrm{Tr}\left[\sigma_\beta \mathcal{E}(\rho_p)\right] = \mathrm{Tr}\left[Q_\beta^\dagger Q_\beta \mathcal{E}(\sum_\alpha p_\alpha \rho_\alpha)\right] = \sum_\alpha \mathrm{Tr}[Q_\beta^\dagger Q_\beta \mathcal{E}(\rho_\alpha)] p_\alpha = \sum_\alpha p_{\beta|\alpha} p_\alpha = f_\beta$. Adding and subtracting $H(f) = -\sum_\beta f_\beta \ln(f_\beta)$ we thus arrive at Eq.M. (11).

Projective measurements: Eq.M. (13)

Next we prove Eq.M. (13a) using a similar technique. Starting from Eq.M. (12) we have

$$\langle v \rangle = \frac{d}{d\lambda} \mathrm{Tr}\left[\rho_q^{-\lambda} \mathcal{E}\left(\rho_p^{\lambda+1}\right)\right] \Big|_{\lambda=0} \tag{5a}$$

$$= \mathrm{Tr}\left[\rho_q^{-\lambda} \mathcal{E}\left(\rho_p^{\lambda+1} \ln(\rho_p)\right) - \rho_q^{-\lambda} \ln(\rho_q) \mathcal{E}\left(\rho_p^{\lambda+1}\right)\right] \Big|_{\lambda=0} \tag{5b}$$

$$= \mathrm{Tr}\left[\mathcal{E}\left(\rho_p \ln(\rho_p)\right)\right] - \mathrm{Tr}\left[\ln(\rho_q) \mathcal{E}\left(\rho_p\right)\right] \tag{5c}$$

$$= S\left(\mathcal{E}(\rho_p) \| \rho_q\right) + S\left(\mathcal{E}\left(\rho_p\right)\right) - S\left(\rho_p\right) \; , \tag{5d}$$

where to arrive at Eq. (5d) we used the fact that $\mathcal{E}$ is trace-preserving, and added and subtracted $S\left(\mathcal{E}\left(\rho_p\right)\right)$.

Finally, Eq.M. (13b) amounts to the following calculation, starting from Eq. (5c):

$$\mathrm{Tr}\left[\mathcal{E}\left(\rho_p \ln(\rho_p)\right)\right] - \mathrm{Tr}\left[\ln(\rho_q)\mathcal{E}\left(\rho_p\right)\right] \tag{6a}$$

$$= S(\rho_q) + \mathrm{Tr}\left[\rho_p \ln(\rho_p)\right] - \mathrm{Tr}\left[\ln(\rho_q)\mathcal{E}\left(\rho_p\right)\right] - S(\rho_q) \tag{6b}$$

$$= S(\rho_q) - S(\rho_p) + \mathrm{Tr}[(\rho_q - \mathcal{E}(\rho_p)) \ln \rho_q] \ . \tag{6c}$$

The heat term

In the main text we claimed that in the thermal case, when $\rho_q = e^{-\beta H_f}/Z_f$ (where H_f is the final Hamiltonian), we find $(2') := \mathrm{Tr}[(\rho_q - \mathcal{E}(\rho_p))\ln(\rho_q)] = -\beta Q$. Here is the proof:

$$\mathrm{Tr}[(\rho_q - \mathcal{E}(\rho_p))\ln(\rho_q)] \tag{7a}$$

$$= -\mathrm{Tr}[(\rho_q - \mathcal{E}(\rho_p))(\beta H_f + \ln Z_f)] \tag{7b}$$

$$= \beta(\mathrm{Tr}[H_f \mathcal{E}(\rho_p)] - \mathrm{Tr}[H_f \rho_q]) \tag{7c}$$

$$= \beta(\langle H_f \rangle_{\mathcal{E}(\rho_p)} - \langle H_f \rangle_{\rho_q}) \ , \tag{7d}$$

where to arrive at Eq. (7c) we used $\mathrm{Tr}[\mathcal{E}(\rho_p)\ln Z_f] = \ln Z_f$ since $\mathcal{E}$ is trace-preserving, and $\mathrm{Tr}[\rho_q \ln Z_f] = \ln Z_f$. Since the (virtual) relaxation process from the state $\mathcal{E}(\rho_p)$ to the state ρ_q is undriven (i.e., there is no work involved), the internal energy change expressed in Eq. (7d) is a pure heat exchange.

DERIVATION OF EQ.M. (17)

Using Eq.M. (9) and Eq.M. (14b), we have

$$\chi_{\mathcal{E}}(\lambda) := \int_{-\infty}^{\infty} dv P_{\mathcal{E}}(v) e^{\lambda v} \tag{8a}$$

$$= \int_{-\infty}^{\infty} dv e^{\lambda v} \sum_{\alpha,j,\beta} \delta(v - V_{\alpha j \beta}) \mathrm{Tr}\left[Q_{\beta|j}\mathcal{E}_j(P_\alpha)\right] p_\alpha \tag{8b}$$

$$= \sum_{\alpha,j,\beta} (p_\alpha/q_{\beta|j})^\lambda \mathrm{Tr}\left[Q_{\beta|j}\mathcal{E}_j(P_\alpha)\right] p_\alpha \tag{8c}$$

$$= \sum_j \mathrm{Tr}\left[\mathcal{E}_j\left(\sum_\alpha p_\alpha^{\lambda+1} P_\alpha\right) \sum_\beta q_{\beta|j}^{-\lambda} Q_{\beta|j}\right] \tag{8d}$$

$$= \sum_j \mathrm{Tr}\left[\mathcal{E}_j\left(\rho_p^{\lambda+1}\right) \rho_{q|j}^{-\lambda}\right] \ , \tag{8e}$$

so that

$$\chi_{\mathcal{E}}(\lambda - 1) = \sum_j \mathrm{Tr}\left[\mathcal{E}_j\left(\rho_p^{\lambda}\right) \rho_{q|j}^{1-\lambda}\right] \tag{9a}$$

$$= \sum_j \mathrm{Tr}\left[\rho_p^\lambda \mathcal{E}_j^*\left(\rho_{q|j}^{1-\lambda}\right)\right] \ . \tag{9b}$$

On the other hand, using Eq.M. (9) and Eq.M. (15b), we have:

$$\tilde{\chi}_{\mathcal{E}^*}(-\lambda) = \int_{-\infty}^{\infty} dv \tilde{P}_{\mathcal{E}^*}(v) e^{-\lambda v} \tag{10a}$$

$$= \int_{-\infty}^{\infty} dv \sum_{\alpha,j,\beta} \delta(v - \tilde{V}_{\beta j \alpha}) \mathrm{Tr}\left[P_\alpha \mathcal{E}_j^*\left(Q_{\beta|j}\right)\right] q_{\beta|j} \tag{10b}$$

$$= \sum_{\alpha,j,\beta} (p_\alpha/q_{\beta|j})^\lambda \mathrm{Tr}\left[P_\alpha \mathcal{E}_j^*\left(Q_{\beta|j}\right)\right] q_{\beta|j} \tag{10c}$$

$$= \sum_j \mathrm{Tr}\left[\sum_\alpha p_\alpha^\lambda P_\alpha \mathcal{E}_j^*\left(\sum_\beta q_{\beta|j}^{1-\lambda} Q_{\beta|j}\right)\right] \tag{10d}$$

$$= \sum_j \mathrm{Tr}\left[\rho_p^\lambda \mathcal{E}_j^*\left(\rho_{q|j}^{1-\lambda}\right)\right] \tag{10e}$$

$$= \chi_{\mathcal{E}}(\lambda - 1) \ , \tag{10f}$$

which is the result in Eq.M. (17).

RECOVERING KNOWN FLUCTUATION THEOREMS

In order to recover the well-established closed system results [4], we consider the CPTP map $\mathcal{E}$ to be simply the unitary evolution:

$$\mathcal{E}(X) = U(t_f, 0) X U^\dagger(t_f, 0) \ . \tag{11}$$

Since this map is unital, its dual is also a CPTP map (the actual time reversed process) given by:

$$\mathcal{E}^*(X) = U^\dagger(t_f, 0) X U(t_f, 0) \ , \tag{12}$$

where $U(t_f, 0) = T_+ \exp(-i \int_0^{t_f} H(t) dt)$, and the Hamiltonian $H(t)$ has instantaneous eigenenergies $\varepsilon(t)$. Using Eq.M. (6) this yields $\gamma = 1$.

Recall that in our formalism we need to also specify the fiducial initial state ρ, the measurements $\mathcal{P}$ and $\mathcal{Q}$, and the distribution q. We pick these so that they generate the Gibbs distributions

$$p_\alpha = \frac{e^{-\beta \varepsilon_\alpha(0)}}{Z(0)} \ , \quad q_\beta = \frac{e^{-\beta \varepsilon_\beta(t_f)}}{Z(t_f)} \tag{13}$$

at inverse temperature β, where $Z(t) = \mathrm{Tr}[\exp(-\beta H(t)]$ is the partition function corresponding to $H(t)$, and $\rho_{\mathrm{G}}(t) = \exp(-\beta H(t))/Z(t)$ is the corresponding Gibbs state. For example, we can assume that $\rho = \rho_{\mathrm{G}}(0)$, $\mathcal{P} = \{|\varepsilon_\alpha(0)\rangle\langle\varepsilon_\alpha(0)|$, and $\mathcal{Q} = \{|\varepsilon_\beta(t_f)\rangle\langle\varepsilon_\beta(t_f)|\}$.

If we let $V_{\{\alpha,\beta\}} = \ln(p_\alpha/q_\beta)$ this then corresponds to the choice:

$$V_{\{\alpha,\beta\}} = \beta(\varepsilon_\beta(t_f) - \varepsilon_\alpha(0)) - \beta(F(t_f) - F(0)) \ , \tag{14}$$

where F is the free energy given by $F(t) = -\ln Z(t)/\beta$. This corresponds to identifying v with the (dimensionless) work.



FIG. 4. The feedback protocol associated with the forward process with an intermediate measurement whose result introduces the conditional CP map $\mathcal{E}_j$.

We thus find from Eq.M. (7) the QJE for a closed quantum system:

$$\langle e^{-\beta(\Delta E-\Delta F)}\rangle = 1 \ , \tag{15}$$

where ΔE is a random variable taking values in the set $\{\varepsilon_\beta(t_f)-\varepsilon_\alpha(0)\}$, and $\Delta F := F(t_f)-F(0)$. For these choices, Eq.M. (13a) becomes

$$\beta\left(\langle\Delta E\rangle - \Delta F\right) = S\left(\mathcal{E}(\rho_{\mathrm{G}}(0))\|\rho_{\mathrm{G}}(t_f)\right) \ , \tag{16}$$

where we have used $S(\mathcal{E}(\rho_{\mathrm{G}}(0)) = S\left(\rho_{\mathrm{G}}(0)\right)$. For a closed quantum system, the heat transfer in and out of the system is zero, so ΔE is equal to the work w. Furthermore, for a closed quantum system, the relative entropy $S\left(\mathcal{E}(\rho_{\mathrm{G}}(0))\|\rho_{\mathrm{G}}(t_f)\right)$ is equal to the mean irreversible entropy production $\langle\Delta S_{\mathrm{irr}}\rangle$ [17, 31], so we can rewrite Eq. (16) as

$$\beta\left(\langle w\rangle - \Delta F\right) = \langle\Delta S_{\mathrm{irr}}\rangle \ , \tag{17}$$

which is the 2nd law of thermodynamics, since $\langle\Delta S_{\mathrm{irr}}\rangle \geq 0$.

As this example illustrates, our formalism clarifies the subtle relationship between the choice of the thermodynamic observable $V_{\{\alpha,\beta\}}$, the initial state probability p_α, and the dual reverse state probability q_β, which together comprise the Jarzynski equality and lead to the 2nd law.

RECOVERING KNOWN FEEDBACK CONTROL RESULTS

Our generalized feedback control protocol is illustrated in Fig. 4.

We can recover known feedback control cases [2, 20] as follows. Assume that the evolution is unitary and the intermediate measurement is error-free. In this case, the jth CP map is given by

$$\mathcal{E}_j(X) = U_jQ_jUXU^\dagger Q_jU_j^\dagger \ . \tag{18}$$

This leads via Eq.M. (16) to

$$\begin{aligned}\gamma &= \sum_j \mathrm{Tr}\left[\rho_{q|j}\mathcal{E}_j(\mathbb{1})\right] = \sum_j \mathrm{Tr}\left[\rho_{q|j}U_jQ_jU_j^\dagger\right]\\ &= \sum_j \mathrm{Tr}\left[Q_jU_j^\dagger\rho_{q|j}U_jQ_j\right] \ .\end{aligned} \tag{19}$$

If we now pick the fiducial initial state ρ, the measurements $\mathcal{P}$ and $\mathcal{Q}'_j$, and the distribution q_j so that they generate the Gibbs distributions

$$p_\alpha = \frac{e^{-\beta\varepsilon_\alpha}}{Z(0)} \ , \quad q_{\beta|j} = \frac{e^{-\beta\varepsilon^j_\beta}}{Z_j(t_f)} \ , \tag{20}$$

where $\rho_{q|j} = \exp(-\beta H_j(t_f))/Z_j(t_f)$ is the Gibbs state associated with $H_j(t_f)$, then we recover the result for γ derived in Ref. [20].

We next consider the case of a classical measurement error. In this scenario when a projection on j is made, there is a measurement error that gives j' with probability $p_{j'|j}$, so that the feedback operation applies $U_{j'}$ instead of U_j. The CP map is then given by:

$$\mathcal{E}_{j'}(X) = \sum_j p_{j'|j}U_{j'}Q_jUXU^\dagger Q_jU_{j'}^\dagger \ . \tag{21}$$

Choosing the same p and q distributions as in Eq. (20) with j replaced by j', results in

$$\begin{aligned}\gamma &= \sum_{j'} \mathrm{Tr}\left[\rho_{q|j'}\mathcal{E}_{j'}(\mathbb{1})\right] = \sum_{j,j'} p_{j'|j}\mathrm{Tr}\left[\rho_{q|j'}U_{j'}Q_jU_{j'}^\dagger\right]\\ &= \sum_{j,j'} p_{j'|j}\mathrm{Tr}\left[Q_jU_{j'}^\dagger\rho_{q|j'}U_{j'}Q_j\right] \ ,\end{aligned} \tag{22}$$

which is again exactly the result for γ derived in Ref. [20].

We are free to choose a different generalized thermodynamic observable:

$$V_{\alpha jj'\beta} = \ln\left(p_\alpha/q_{\beta|j'}\right) + I_{jj'} =: -\tilde{V}_{\beta j'j\alpha} \ , \tag{23}$$

where

$$I_{jj'} := \ln\left(p_{j'|j}/p_{j'}\right) \ , \tag{24}$$

the mean of which is the classical mutual information. Note that

$$\sum_{\beta,j'} p_{j'}q_{\beta|j'} = 1 \ . \tag{25}$$

The joint distribution can be decomposed using Bayes' rule as $p_{(\alpha,j,j',\beta)} = p_\alpha p_{(j,j',\beta)|\alpha}$, where the conditional probability is $p_{(j,j',\beta)|\alpha} = p_{j'|j}\mathrm{Tr}\left[Q_{\beta|j'}U_{j'}Q_jUP_\alpha U^\dagger Q_jU_{j'}^\dagger\right]$.

We then have

$$P_{\{\mathcal{E}_{j'}\}}(v) = \sum_{\alpha,j,j'\beta} \delta(v-V_{\alpha jj'\beta})p_{(\alpha,j,j',\beta)} \tag{26a}$$

$$\begin{aligned}&= \sum_{\alpha,j,j'\beta} \delta(v-V_{\alpha jj'\beta})p_\alpha p_{j'|j}\\ &\quad\times \mathrm{Tr}\left[U^\dagger Q_jU_{j'}^\dagger Q_{\beta|j'}U_{j'}Q_jUP_\alpha\right] \ ,\end{aligned} \tag{26b}$$

so that

$$P_{\{\mathcal{E}_{j'}\}}(v)e^{-v} = \sum_{\alpha,j,j'\beta} \delta(-v - \tilde{V}_{\beta j' j\alpha}) q_{\beta|j'} p_{j'} \times \mathrm{Tr}\left[U^\dagger Q_j U_{j'}^\dagger Q_{\beta|j'} U_{j'} Q_j U P_\alpha\right] \ . \quad (27)$$

Note that the quantity

$$\sum_{\beta,j'} p_{j'} q_{\beta|j'} U_{j'}^\dagger Q_{\beta|j'} U_{j'} =: \hat{\rho} \quad (28)$$

is a density matrix. We can define a CPTP map:

$$\hat{\mathcal{E}}(\hat{\rho}) = U^\dagger \left[\sum_j Q_j \hat{\rho} Q_j\right] U \ . \quad (29)$$

We thus recover the integrated fluctuation theorem:

$$\int dv P_{\mathcal{E}}(v) e^{-v} = \sum_\alpha \mathrm{Tr}\left[P_\alpha \hat{\mathcal{E}}(\hat{\rho})\right] = \mathrm{Tr}\left[\hat{\mathcal{E}}(\hat{\rho})\right] = 1 \ , \quad (30)$$

which is again the result in Ref. [20].

DETAILS ABOUT THE MASTER EQUATION

The underlying model assumes a total Hamiltonian of the form $H(t) = H_S(t) + H_B + H_I$, where H_B is the bath Hamiltonian and H_I is the system-bath interaction. We consider a simple harmonic oscillator bath $H_B = \sum_k \omega_k b_k^\dagger b_k$, with the interaction given by the standard dephasing model [54], $H_I = g \sum_i \sigma_i^z \otimes \sum_k \left(b_k + b_k^\dagger\right)$.

The master equation used in our simulations is derived in Ref. [44] and is given by (in units of $\hbar = 1$) :

$$\dot{\rho}_S(t) = -i\left[H_S(t) + H_{\mathrm{LS}}(t), \rho_S(t)\right] + \sum_\omega \gamma(\omega) \times \left(L_\omega(t)\rho_S(t)L_\omega^\dagger(t) - \frac{1}{2}\left\{L_\omega^\dagger(t)L_\omega(t), \rho_S(t)\right\}\right) \quad (31)$$

where H_{LS} is the Lamb shift, the γ's are dephasing and relaxation/excitation rates. The Lindblad operators are given by:

$$L_\omega(t) = \sum_{\omega=\varepsilon_b(t)-\varepsilon_a(t)} |\varepsilon_a(t)\rangle\langle\varepsilon_a(t)| \sum_i \sigma_i^z |\varepsilon_b(t)\rangle\langle\varepsilon_b(t)| \ , \quad (32)$$

where the instantaneous Bohr frequency ω is expressed in terms of the instantaneous energy eigenstates, i.e., $H_S(t)|\varepsilon_a(t)\rangle = \varepsilon_a(t)|\varepsilon_a(t)\rangle$ For an Ohmic bath with high-frequency cut-off ω_c, we have

$$\gamma(\omega) = \frac{g^2 \eta \omega e^{-\omega/\omega_c}}{1 - e^{-\beta\omega}} \ , \quad H_{\mathrm{LS}} = \sum_\omega S(\omega) L_\omega^\dagger(t) L_\omega(t) \ , \quad S(\omega) = \int_{-\infty}^{\infty} \frac{d\omega'}{2\pi} \gamma(\omega') \mathcal{P}\left(\frac{1}{\omega - \omega'}\right) \ . \quad (33)$$



FIG. 5. Independent calculation of $\langle e^{-\beta(\Delta E - \Delta F)}\rangle$ (blue solid) and γ (red dashed) as a function of the system-bath coupling strength using the adiabatic Markovian master equation with $t_f = 5\mu s$, $J = 1/2$, $T = 17$mK, and $\omega_c = 8\pi$. Inset: magnification of the behavior near the top of the curves. The two curves overlap to within numerical accuracy.

where η is a parameter (with dimension time squared) characterizing the Ohmic bath, and $\mathcal{P}$ denotes the Cauchy principal value. To see that this CP map is not unital, we plug $\mathbb{1}$ into the RHS of Eq. (31) and find that the non-zero component arises from the dissipative part associated with relaxation and excitation processes

$$\sum_\omega \gamma(\omega) \left[L_\omega(t), L_\omega^\dagger(t)\right] \ . \quad (34)$$

Since $\gamma(\omega) \neq \gamma(-\omega)$, meaning that the relaxation and excitation transition rates are unequal, the term (34) is non-zero, making the CPTP map generated by the master equation non-unital.

NUMERICAL CONFIRMATION OF THE QJE AND FIRST MOMENT EXPRESSION FOR THE ADIABATIC MARKOVIAN MASTER EQUATION

In order to test Eq.M. (18) and Eq.M. (13a) we performed the following simulations for the two-qubit model described in the main text. We initialized the system in one of the four energy eigenstates, $\rho_S(0) = |\varepsilon_a(0)\rangle\langle\varepsilon_a(0)|$, then we evolved the density matrix using our adiabatic master equation. The diagonal elements of the density matrix at $t = t_f$ are then associated with the probability $p_{\beta|\alpha}$ of measuring the state $|\varepsilon_\beta(t_f)\rangle$. Using this we calculated the expectation value $\langle e^{-\beta(\Delta E - \Delta F)}\rangle$. We then used our adiabatic master equation to evolve the identity operator. This allowed us to numerically find $\mathcal{E}(\mathbb{1})$, which in turn allowed us to calculate γ. The equality expressed in Eq.M. (18) is obtained with high accuracy, as shown in Fig. 5. We can also calculate the LHS and RHS of the first moment of our fluctuation theorem [Eq.M. (13a)] independently. We again find excellent agreement between the two results: see Fig. 6.



FIG. 6. Simulation results for $\langle \beta (\Delta E - \Delta F) \rangle$ (solid blue) and the RHS of Eq. (13a) (red dashed) using the Markovian master equation and $t_f = 5\mu s$, $J = 1/2$, $T = 17$mK, and $\omega_c = 8\pi$. The two curves overlap to within numerical accuracy.

To compute ΔF, as needed for Eq.M. (20), the eigenvalues ε_i of the initial ($i = \alpha$) and final ($i = \beta$) Hamiltonian are numerically computed by diagonalizing the respective Hamiltonians. In turn, the respective partition functions are calculated using the energy eigenvalues found, $Z = \sum_i e^{-\beta\varepsilon_i}$. The free energy is then $F = -\ln(Z)/\beta$.

EXPERIMENTAL SYSTEM

Our experiments were performed using the D-Wave One Rainier chip at the USC Information Sciences Institute, comprising 16 unit cells of 8 superconducting flux qubits each, with a total of 108 functional qubits. The couplings are programmable superconducting inductances. The qubits and unit cell, readout, and control have been described in detail elsewhere [55–57]. The initial energy scale for the transverse field is 33.7GHz (the A function in Fig 7), the final energy scale for the Ising Hamiltonian (the B function) is 33.6GHz, about 15 times the experimental temperature of 17mK $\approx$ 2.3GHz. For a given Ising Hamiltonian, the quantum annealing process was repeated 50,000 times per qubit pair. We used five different qubit pairs to rule out systematic local magnetic field bias. As a further precaution against systematic bias we applied three spin-inversion transformations to our Ising Hamiltonian: $H_{\text{Ising}} \mapsto \sigma_j^x H_{\text{Ising}} \sigma_j^x$ for $j = 1, 2$, and $H_{\text{Ising}} \mapsto \sigma_1^x \sigma_2^x H_{\text{Ising}} \sigma_2^x \sigma_1^x$, all of which commute with the transverse field component $\sum_i \sigma_i^x$ of our system Hamiltonian. These transformations simply relabel the energy spectrum, i.e., if a certain spin configuration has energy E, then under the transformation $\sigma_j^x H_{\text{Ising}} \sigma_j^x$, the configuration with the j-th spin flipped will also have energy E. Averaging the results over the four different isospectral Ising Hamiltonians and over the five different qubit pairs for a given Ising problem, we have a total of 10^6 data points per given values of J and t_f.

At $J = 0$, we would expect to find the (excited) states $|\uparrow\downarrow\rangle$ and $|\downarrow\uparrow\rangle$ with equal probability. However, this symmetry is broken in our experimental data. This should not be interpreted as being solely due to a local magnetic field bias, since averaging over spin-inversion transformations should have cancelled any such bias. This suggests a more systematic (unaccounted) source of noise in the experiment. We found that to fit the experimental data in Fig.M. 3 over the full range of Ising couplings J, the local magnetic fields in the Ising Hamiltonian needed to be adjusted from their ideal value of 1/3 to $h_1 = 0.28$ and $h_2 = 0.37$.

The theoretical best fit (in Fig.M. 3) was found by determining the value of $\kappa = g^2\eta/\hbar^2$ that minimizes the mean square deviation (MSD) between the n experimental $\{\langle v\rangle_{\text{Ex},i}\}$ and theoretical $\{\langle v\rangle_{\text{Th},i}\}$ results:

$$\text{MSD}(\kappa) = \frac{1}{n}\sum_{i=1}^{n} \left(\langle v\rangle_{\text{Ex},i} - \langle v(\kappa)\rangle_{\text{Th},i}\right)^2 \quad . \tag{35}$$

In principle, the high-frequency cut-off ω_c is also a free parameter in our theoretical model, and it can also be used as part of the fitting parameters. We found that choosing a different ω_c requires a different optimal κ value to fit the data, but we restricted ourselves to $\omega_c = 8\pi$ GHz since it nicely satisfies the approximations made in the derivation of the master equation [44].



FIG. 7. The annealing schedules $A(t)$ and $B(t)$ used in the system Hamiltonian in the experiment and in our numerical simulations.